

PCT/EP+ 00/08157




Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

EP 00/08157



Bescheinigung

Certificate

4
10/069575
Attestation

Die angehefteten Unterlagen stimmen mit der ursprünglich eingereichten Fassung der auf dem nächsten Blatt bezeichneten europäischen Patentanmeldung überein.

The attached documents are exact copies of the European patent application described on the following page, as originally filed.

Les documents fixés à cette attestation sont conformes à la version initialement déposée de la demande de brevet européen spécifiée à la page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

99402149.1

PRIORITY DOCUMENT

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

I.L.C. HATTEN-HECKMAN

DEN HAAG, DEN
THE HAGUE, 27/11/00
LA HAYE, LE



Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

Blatt 2 der Bescheinigung
Sheet 2 of the certificate
Page 2 de l'attestation

Anmeldung Nr.:
Application no.: 99402149.1
Demande n°:

Anmeldetag:
Date of filing: 30/08/99
Date de dépôt:

Anmelder:
Applicant(s):
Demandeur(s):
I. D. M. IMMUNO-DESIGNED MOLECULES
75011 Paris
FRANCE

Bezeichnung der Erfindung:
Title of the invention:
Titre de l'invention:
New humanized biomaterials

In Anspruch genommene Priorität(en) / Priority(ies) claimed / Priorité(s) revendiquée(s)

Staat:
State:
Pays:

Tag:
Date:
Date:

Aktenzeichen:
File no.
Numéro de dépôt:

Internationale Patentklassifikation:
International Patent classification:
Classification internationale des brevets:
A61L27/40, A61L27/38

Am Anmeldetag benannte Vertragsstaaten:
Contracting states designated at date of filing: AT/BE/CH/CY/DE/DK/ES/FI/FR/GB/GR/IE/IT/LI/LU/MC/NL/PT/SE/TR
Etats contractants désignés lors du dépôt:

Bemerkungen:
Remarks:
Remarques:

See for original title page 1 of the description

NEW HUMANIZED BIOMATERIALS, A PROCESS FOR THEIR PREPARATION AND THEIR APPLICATIONS

5 The invention relates to new humanized biomaterials, a process for their preparation and their applications.

Tissue repair is needed after severe bone fractures, cartilage loss or general fragilization during ageing.

10 Artificial metallic or even ceramic prostheses are not very well integrated within host tissues and replacement with new surgery is often required after a few years, a major problem of handicap for old people.

Autologous grafts of bones or cartilage tissue is very difficult and costly. New technologies develop porous matrices implanted as scaffold prosthesis.

15 These matrices can eventually be filled with growth factors for the tissue regeneration or even with bone marrow stem cells.

However, fixation of cells or factors in the porous matrices with very prolonged and slow release of growth factors is very difficult to achieve and the ideal cocktail and concentration of factors required is unknown.

20 The aim of the present invention is to provide a homogeneous humanized, bioactive biomaterial (for example porous ceramics) that can be used for implantation purposes and which do not present the long term biocompatibility problems of prior art.

25 Another aim of the invention is to provide a bioactive biomaterial enabling tissue growth (for example bone and cartilage) in its porous space and securing the integration of the grafted biomaterial in the surrounding tissues (viable bones...).

Another aim of the invention is to provide long lasting prostheses, which avoids requirement for replacement of biomaterial prostheses after 10 years, as often needed up to now.

30 These aims are achieved by the invention, which consists in humanized biomaterial comprising a porous biocompatible composite material customized and implanted with monocyte derived cells and preferably with macrophages.

The expression "humanized" means that the porous biomaterial has been colonized with human cells derived from blood monocytes.

35 The expression "biocompatible composite" material designates a material composed of one or several of the following materials proved to be non toxic for human tissues (carbon microfibers, ceramics, calcium phosphates, metal oxides, collagen polymers...).

The expression "porous" means that the biomaterial and preferably the ceramic present pores of about 100 to 2000 microns of diameter.

The expression "customized and implanted" material means that the shape and size of biomaterial is designed specifically for a patient and a site of implantation.

5 The expression "monocyte derived cells" corresponds to human mononuclear cells isolated from blood, enriched in monocytes and cultured at 37° C in appropriate medium, for 5 to 10 days to obtain tissue type macrophages.

10 The monocyte derived cells used in the invention are for instance such as those described in PCT/EP 93/01232, WO 99/13054, EP 96/ 901848.0-2107, WO 97/44441.

In a particular embodiment of the invention, the monocyte derived cells described above, contain exogenous compounds such as drugs, proteins, growth factors of interest.

15 In another embodiment, the monocyte derived cells as described above contain in their cytoplasm exogenous DNA coding for a protein of interest.

20 The substantially irreversible humanization of matrices of biocompatible composite material described in the present invention allows a physiological interaction between the prostheses made of the biomedical composite, grafted and the host cells in the body. These relations with host tissue cells and with the extracellular matrix allow reconstruction of epithelial sheets and growth of a capillary network around the grafted biomaterial by local multiplication and sprouting of endothelial cells.

25 The monocyte derived cells, in particular the macrophages used to humanize *in vitro* the porous material in the invention are particularly adequate to increase integration and *in vivo* lifespan of biocompatible prostheses.

Advantageously, the humanized biomaterial of the invention is homogenous.

30 According to an advantageous embodiment, in the humanized biomaterial of the invention, the biocompatible composite material is chosen among the following materials : microfibres, ceramic materials, metal oxides such as aluminum oxide, calcium phosphate ceramic, glass or carbon fibers, hydroxylapatite, silicon carbide or nitride, collagen polymers or a mixture of these different materials.

35 According to another advantageous embodiment, in the humanized biomaterial of the invention, the human macrophages are liable to be obtained by *ex vivo* differentiation from blood monocytes leading to living macrophages, and are cultured under conditions enabling their penetration and adherence into the biomaterial for instance for several hours at 37°C, with the porous biomaterial, allowing infiltration of the biomaterial and substantially irreversible binding of the living macrophages to

the biomaterial, now humanized with patient's macrophages and ready for implantation.

The expression "substantially irreversible binding" means that macrophages are tightly bound by numerous contacts with the material and cannot be detached under physiological conditions.

The invention also relates to a living body-supporting implant, characterized by the fact that it comprises or consists of the humanized biomaterial according to any one of claims 1 to 3, and is preferably structured under the form of scaffold, tissue-supporting sponges, bone or cartilage.

The expression "living body-supporting implant" designates an implant having for example the physical form and robustness of a bone to be replaced.

The expression "scaffold" designates a physical structure eventually metallic which keeps the biomaterial (ceramics or other) in the appropriate conformation.

The expression "tissue supporting sponges" designates a soft implant formed for instance of collagen which will be humanized with macrophages before insertion in the body.

The invention also relates to the use of a humanized biomaterial of the invention or of a living body-supporting implant of the invention, for the preparation of a tissue implant destined to replace or repair defective tissue, such as defective bone, cartilage, dental tissue, fibrous tissue, fibrocartilaginous supporting tissue.

The expression "fibrous tissue" designates tissues surrounding organs which support this organ and maintain the shape of this body part : they are mainly formed by epithelial sheets.

According to another advantageous embodiment, the invention relates to the use of a humanized biomaterial of the invention or of a living body-supporting implant of the invention, wherein the monocyte derived cells or macrophages are autologous with respect to the tissue to be replaced or repaired, enabling the biomaterial or the living body-supporting implant to be recognized as self.

The expression "implant to be recognized as self" means that it contains cells (i.e. macrophages) from the patient in which it will be grafted and is therefore autologous to the host.

A process for the preparation of a humanized biomaterial of the invention comprises the following steps :

- preparation of the porous biomaterial structured in form of bones, cartilage,
- preparation of macrophages from blood monocytes,

- immersion of the biomaterial in a physiologic solution appropriate for the culture of macrophages which are added afterwards (ex. : phosphate buffered saline, medium such as RPMI, IMDM, AIMV),
- addition of the macrophage to the solution under conditions enabling binding to the biomaterial and particularly for 1 to 20 h. at 37°C, 5 % CO₂ and 5 % air,
- washing of the biomaterial and conservation until use in physiologic medium.

5
10 A process for the preparation of a living body-supporting implant of the invention comprises the following steps :

- preparation of a customized porous implant or scaffold composed of bio-compatible material, according to any one of claims 1 to 3,
- preparation of macrophages from blood monocytes of the patient needing the customized implant of biomaterial,
- co-culture of macrophages and the implant in adequate medium under conditions enabling penetration and adherence to the biomaterial, in particular at 37°C, 5% CO₂ in hydrophobic bags or containers until grafting the implant.

15
20 The invention also relates to the use of the humanized biomaterial or of a living body-supporting implant, which can be implanted in a tissue, for the *in vitro* or *in vivo* or *ex vivo* delivery of factors chosen in the group of chemokines and/or monokines, and/or cytokines and/or growth factors, the factors released being useful for the local attraction of cells required for tissue growth (such as osteoblasts, chondrocytes, fibroblasts, epithelial cells.....) and/or for the neovascularization around the implanted biomaterial, and/or for the release of growth factors sustaining proliferation of cells and/or the growth of new tissues.

25
30 Indeed, macrophages maintain tissue homeostasis through the secretion of at least 80 growth factors or monokines controlling and inducing proliferation of mesenchymal (fibroblasts....), endothelial, chondrocytes, osteoblasts, epithelial cells.. They also secrete enzymes and mediators allowing growth and renewal of the surrounding cells and tissues (see Table 1).

35 The key factors secreted by macrophages supporting tissue integration regeneration and growth of mesenchymal cells are : IGF1 and TGFs, but also PDGF, bFGF, MDGF, GM, CSF, NAF, IL-8, TNF, angiogenin and angiogenic factors. These growth factors allow also the development of all the steps required for angiogenesis, allowing neovascularisation and reconstitution of blood microcapillaries around the grafted biomaterial.

In this aspect, macrophages are synthesizing 10 fold more proteins than monocytes, much more growth factors and less inflammatory mediators.

5

TABLE 1 :

GROWTH FACTORS PROTEINS AND MEDIATORS SUPPORTING
TISSUE HOMEOSTASIS SECRETED BY NATURE MACROPHAGES

10

ENZYMES :

Lyzosymes
Neutral proteases
Plasminogen activator
Collagenase
Elastase
Angiotensin-convertase

Acid hydrolases

Proteases
Lipases
Ribonucleases
Phosphatases
Glycosidases
Sulphatases

Arginase

COMPLEMENT COMPONENTS

C1, 4, 2, 3 and 5
Factors B and D and Properdin
C1 inhibitor
C3b inactivator and β -1H

ENZYME INHIBITORS
(Antiproteases)

α 1-antiprotease
Plasmin inhibitors
 α -2 macroglobulin
Plasminogen activator inhibitors

PROTEINS BINDINGMETABOLIC AND LIPIDS :

Acidic isoferritins

BIOACTIVE LIPIDS :

Arachidonic acid metabolites
Prostaglandins E2, F2 α
Prostacyclin
Thromboxane
Leukotrienes B4, C, D and E
Hydroxy-eicosatetraenoic acids
(including SRS-A)
Platelet activating factors

CYTOKINES, HORMONES,GROWTH FACTORS :

Interleukins 1 α and β
Tumours necrosis factor α
Interferons α and β ₁, &
Interleukin 6, 8, 13, 18

Chemotatic factors for

Neutrophils
Tlymphocytes
Monocytes
Fibroblasts

Heamatopoietic Colony Stimulating
Factors for

Granulocyte-Macrophages (GM-CSF)
Granulocytes (G-CSF)
Macrophages (M-CSF)
Erythropoietin

Growth factors

Fibroblasts growth factor /

Insuline like G.F.

"platelet-derived growth factor"
(PDGF)

Transforming growth factor α and β

Endothelial cell growth factor

Hormones

1 α , 25-Dihydroxyvitamin D3

Transferrin
Transcobalamin II
Fibronectin
Laminin
Lipid transfer protein
Thrombospondin

NUCLEOSIDES AND
METABOLITES :
Thymidine and deoxycytidine
Uracil
Uric acid
Lactic acid
Polyamines nitrines and nitrates
Neopterin

Insulin-like activity, protagandins
Thymosin B4
 β endorphin
Adrenocorticotrophic hormone

CHEMOKINES MIP / RANTE FAMILIES
COAGULATION FACTORS :
Factors III, VIII, and tissue factor
Prothrombin and prothrombinase
Factors IX, X, V and VII

In an advantageous embodiment of the invention, the macrophages migrate initially in the porous biomaterial and incorporate irreversibly into this prosthesis by very strong adherence and spreading. When they are kept in physiological conditions, macrophages are very long living cells lasting from several months to several years after implantation. During this time, macrophages will continuously secrete growth factors and cytokines in their local environment ; these factors will act in synergy at very low concentrations (10^{-10} M) on the surrounding cells and tissues.

Furthermore, macrophages do present on their membranes receptors for cytokines, hormones, sugars allowing to respond to micro-environmental needs and to adjust their secretion to the local status around the biomaterial at different periods after grafting.

The growth factors secreted by macrophages represent the global requirements for tissue repair, differentiation and local angiogenesis. The chemokines which will be continuously released in a concentration gradient around the implant will call in and around the prosthesis cells required for recolonization and integration of the biomaterial in host environment. Therefore, the new customized porous biomaterials colonized with host macrophages present a novel biotolerance and a length of adequate performance far better than prosthesis used in the absence of autologous macrophages. Applications are very large in solid or cartilaginous prosthesis needed in bone, cartilaginous repair particularly.

According to another advantageous embodiment, the invention relates to the use of the humanized biomaterial or of a living body-supporting implant, as a graft for the replacement of supporting tissues such as bones, cartilages, dental tissues, epithelial sheet and subcutaneous tissue matrix.

Example 1 :

5 A calcium phosphate porous ceramic with pores of 200 to 2000 microns (porosity > 20 % and < 80 %) is placed on and hydrophobic support (ethylene vinyl acetate) in the presence of 50 ml AIMV culture medium (life-cell Gibco, Paisley G.B.). Macrophages are added to this preparation at the concentration of $5 \cdot 10^6/\text{ml}$; they are obtained after 7 days differentiation of blood monocytes in culture according to published state of the art in publications and patents (PCT/EP 93/01232, WO 99/13054, EP 96/ 901848.0-2107, WO 97/44441) ; a control preparation is
10 maintained in the absence of macrophages. The preparation is incubated overnight at 37 °C, 5 % CO₂, 95 % air to allow fixation of macrophages on the ceramic.

The porous ceramic is washed and cultured in the presence of fibroblasts and/or in the presence of chondrocytes. In both cases, the cell proliferation is higher by a factor 2 to 10 for the porous ceramic colonized with macrophages, compared to
15 control ceramic.

Example 2 :

20 A small fragment of porous microceramic is implanted in a rabbit cornea. The inert microceramic piece induces a very small to moderate and only peripheral growth of new blood vessels from the ring of the cornea.

In contrast, microceramic covered with macrophages, as described in example 1 induces a major neovascularisation towards the center of the cornea. The cornea becomes vascularized through an invasion of endothelial cells arising from the rim
25 rich in blood supply and sprouting towards the biomaterial implanted.

CLAIMS

5 1 - Humanized biomaterial comprising a porous biocompatible composite material customized and implanted with monocyte derived cells and preferably with macrophages.

10 2 - Humanized biomaterial according to claim 1, wherein the biocompatible composite material is chosen among the following materials : microfibers, ceramic materials, metal oxides such as aluminum oxide, calcium phosphate ceramic, glass or carbon fibers, hydroxylapatite, silicon carbide or nitride, collagen polymers or a mixture of these different materials.

15 3 - Humanized biomaterial according to claims 1 or 2, wherein the human macrophages are liable to be obtained by *ex vivo* differentiation from blood monocytes leading to living macrophages, and are cultured under conditions enabling their penetration and adherence into the biomaterial, for instance for several hours at 37°C, with the porous biomaterial, allowing infiltration of the biomaterial and substantially
20 irreversible binding of the living macrophages to the biomaterial, being humanized with patient's macrophages and ready for implantation.

25 4 - Living body-supporting implant, characterized by the fact that it comprises or consists of the humanized biomaterial according to any one of claims 1 to 3, and is preferably structured under the form of scaffold, tissue-supporting sponges, bone or cartilage.

30 5 - Use of a humanized biomaterial according to any one claims 1 to 3 or of a living body-supporting implant according to claim 4, for the preparation of a tissue implant destined to replace or repair defective tissue, such as defective bone, cartilage, dental tissue, fibrous tissue, fibrocartilaginous supporting tissue.

35 6 - Use of a humanized biomaterial according to any one of claims 1 to 3 or of a living body-supporting implant according to claim 4, wherein the monocyte derived cells or macrophages are autologous with respect to the tissue to be replaced or repaired, enabling the biomaterial or the living body-supporting implant to be recognized as self.

7 - Process for the preparation of a humanized biomaterial according to any one claims 1 to 3, comprising the following steps :

- 5 - preparation of the porous biomaterial structured in form of bones, cartilage,
- preparation of macrophages from blood monocytes,
- immersion of the biomaterial in a physiologic solution appropriate for the culture of macrophages which are added afterwards (ex. : phosphate buffered saline, medium such as RPMI, IMDM, AIMV),
- 10 - addition of the macrophages to the solution under conditions enabling binding to biomaterial and particularly for 1 to 20 h. at 37°C,
- 5 % CO₂ and 5 % air,
- washing of the biomaterial and conservation until use in physiologic medium.
- 15

8 - Process for the preparation of a living body-supporting implant according to claim 4, comprising the following steps :

- 20 - preparation of a customized porous implant or scaffold composed of bio-compatible material, according to any one of claims 1 to 3,
- preparation of macrophages from blood monocytes of the patient needing the customized implant of biomaterial,
- co-culture of macrophages and the implant in adequate medium
- 25 under conditions enabling penetration and adherence to the biomaterial in particular at 37°C, 5% CO₂ in hydrophobic bags or containers until grafting the implant.

9 - Use of the humanized biomaterial according to any one of claims 1 to 3 or of a living body-supporting implant according to claim 4, which can be implanted in a tissue, for the *in vitro* or *in vivo* or *ex vivo* delivery of factors chosen in the group of chemokines and/or monokines, and/or cytokines and/or growth factors, the factors released being useful for the local attraction of cells required for tissue growth (such as osteoblasts, chondrocytes, fibroblasts, epithelial cells.....) and/or for the

30

35 neovascularization around the implanted biomaterial, and/or for the release of growth factors sustaining proliferation of cells and/or the growth of new tissues.

10 - Use of the humanized biomaterial according to any one of claims 1 to 3 or of a living body-supporting implant according to claim 4, as a graft for the replacement of supporting tissues such as bones, cartilages, dental tissues, epithelial sheet and subcutaneous tissue matrix.

ABSTRACT**NEW HUMANIZED BIOMATERIALS, A PROCESS FOR THEIR
PREPARATION AND THEIR APPLICATIONS**

5

The invention relates to humanized biomaterial comprising a porous biocompatible composite material customized and implanted with monocyte derived cells preferably with macrophages

10

(no figure)

